

Salt Tolerance of Methicillin-Resistant and Methicillin-Susceptible *Staphylococcus aureus* ▽

The prevalence of methicillin-resistant *Staphylococcus aureus* (MRSA) strains causing nosocomial infections is still increasing worldwide, resulting in high morbidity and mortality rates and increased hospital costs (5, 6). So far, in hospitals in The Netherlands, a strict search-and-destroy policy, including screening for MRSA carriage by culturing, has kept the prevalence of MRSA under 1% (6, 8).

To increase the sensitivity of culturing methods, a mannitol salt enrichment broth is commonly used. This medium traditionally contains 7.5% sodium chloride as a selective agent based on the long-known high salt tolerance of staphylococci (1, 4). The *Clinical Microbiology Procedures Handbook*, published by the American Society for Microbiology, recommends using a broth with 6.5% or 7.5% NaCl (2).

In 1997, however, Jones et al. found that this salt enrichment broth inhibited the growth of the epidemic strain MRSA-16 (EMRSA-16) (3). They determined that the MIC of NaCl for EMRSA-16 was 7% and that low inocula of EMRSA-16 were inhibited by NaCl concentrations higher than 2.5%.

To optimize the MRSA screening protocol used in our laboratory, we determined the salt tolerance of a well-defined collection of 38 methicillin-resistant *S. aureus* strains (*mecA* gene positive with oxacillin MICs of 0.38 to ≥ 256 $\mu\text{g/ml}$ and various phage types) and 45 methicillin-susceptible *S. aureus* (MSSA) strains (*mecA* gene negative with oxacillin MICs of 0.25 to 12 $\mu\text{g/ml}$).

We prepared a mannitol base broth to which we added NaCl to final concentrations ranging from 1.5 to 12.5%, with intervals of 1%. The mannitol salt broths were dispensed in microdilution trays, and for each strain, wells with concentrations of NaCl increasing from 1.5 to 12.5% were inoculated with a suspension to a final concentration of 10^2 CFU/ml. After 24 and 48 h of incubation, we read the NaCl maximal noninhibiting concentrations (MaNICs) and NaCl MICs for each strain.

After 24 h, 11 (29%) MRSA strains of different phage types had MaNICs of 2.5% (Fig. 1), meaning that for roughly one-third of the tested MRSA strains, growth was inhibited by an NaCl concentration higher than 2.5%. Figure 2 shows the NaCl MICs for MRSA. After 24 h, the MIC₉₀ was 6.5%,




FIG. 1. NaCl MaNICs for MRSA strains after 24 and 48 h of incubation.




FIG. 2. NaCl MIC values for MRSA after 24 and 48 h of incubation.

which is lower than the commonly used concentration of NaCl of 7.5%.

For MSSA strains, the MaNICs and MICs were similar to those for MRSA strains. We found no correlation between the oxacillin MIC and the NaCl MaNICs or MICs for either MRSA or MSSA.

Our findings confirm that for optimal sensitivity, an enrichment broth for MRSA screening should not contain more than 2.5% NaCl. This concentration, however, is too low to inhibit contaminating flora sufficiently (7). Thus, for adequate selectivity, antimicrobial agents should be used instead of NaCl. Guidelines for MRSA screening should be adapted to this effect.

We thank Wim J.B. Wannet (RIVM, Bilthoven, The Netherlands) for providing MSSA strains.

REFERENCES

- Chapman, G. H. 1945. The significance of sodium chloride in studies of staphylococci. *J. Bacteriol.* **50**:201–203.
- Isenberg, H. D. (ed.). 2004. *Clinical microbiology procedures handbook*, 2nd ed. American Society for Microbiology, Washington, DC.
- Jones, E. M., K. E. Bowker, R. Cooke, R. J. Marshall, D. S. Reeves, and A. P. MacGowan. 1997. Salt tolerance of EMRSA-16 and its effect on the sensitivity of screening cultures. *J. Hosp. Infect.* **35**:59–62.
- Koch, F. E. 1942. Electivnährboden für Staphylokokken. *Zentr. Bakt. Parasitenk. I Orig.* **149**:122–124.
- National Nosocomial Infections Surveillance System. 2004. National Nosocomial Infections Surveillance (NNIS) System Report, data summary from January 1992 through June 2004, issued October 2004. *Am. J. Infect. Control* **32**:470–485.
- Tiemersma, E. W., S. L. Bronzwaer, O. Lyytikäinen, J. E. Degener, P. Schrijnemakers, N. Bruinsma, J. Monen, W. Witte, and H. Grundman. 2004. Methicillin-resistant *Staphylococcus aureus* in Europe, 1999–2002. *Emerg. Infect. Dis.* **10**:1627–1634.

7. **van Enk, R. A., and K. D. Thompson.** 1992. Use of a primary isolation medium for recovery of methicillin-resistant *Staphylococcus aureus*. *J. Clin. Microbiol.* **30**:504–505.
8. **Verhoef, J., D. Beaujean, H. Blok, A. Baars, A. Meyler, C. van der Werken, and A. Weersink.** 1999. A Dutch approach to methicillin-resistant *Staphylococcus aureus*. *Eur. J. Clin. Microbiol. Infect. Dis.* **18**:461–466.

Marjan J. Bruins*

Petra Juffer

Maurice J. H. M. Wolfhagen

Gijs J. H. M. Ruijs

Laboratory of Clinical Microbiology and Infectious Diseases

Isala klinieken

Rhijnvis Feithlaan 62

8021 AM Zwolle, The Netherlands

*Phone: 31384243111

Fax: 31384243146

E-mail: m.j.bruins@isala.nl

[†] Published ahead of print on 13 December 2006.